

Characteristics

ADOB[®] 2.0 Mn is a high-purity, oligo-nutrient fertiliser containing 10.1% manganese (Mn), 6.5% nitrate-nitrogen (N-NO₃) and 2% magnesium (MgO). It is a homogeneous aqueous solution, which makes it user-friendly and ensures that nutrients are highly available to plants' leaves and roots. It can be applied through fertigation, hydroponics and foliar spray. It is especially suited to foliar application since the unique production "Technology 2.0" greatly enhances foliar uptake of manganese.

ADOB[®] 2.0 Mn controls and prevents manganese deficiency in all arable and horticultural crops and under all management methods, even in highly challenging conditions. Correction of manganese deficiency amends the physiological disorders connected with the insufficient activity of many enzymes which catalyse redox, decarboxylation and hydrolytic reactions. This in turn increases oil concentration in seeds and boosts plant resistant to root-infecting pathogens, directly boosting crop performance.







preventive and corrective effect



Packaging: 10, 20, 1000 |

Composition

Composition – ADOB[®] 2.0 Mn

Nutrients	Symbol	Content [% w/w]	Content [% w/v]	Content [g/l]	Form
Total nitrogen	Ν	6.5	9.1	91.0	
– nitrate nitrogen	N-NO₃	6.5	9.1	91.0	
Magnesium oxide	MgO	2.0	2.8	28.0	soluble in water
Manganese	Mn	10.1	14.1	141.0	soluble in water



Przedsiębiorstwo Produkcyjno-Consultingowe ADOB Sp. z o.o. Sp. k.

ul. Kołodzieja 11 61-070 Poznań, PL email: office@adob.com.pl





Application recommendations

Application recommendations – ADOB[®] 2.0 Mn

	Crops	Number of applications per season	Crop phenological stage	BBCH stage	Product application rate [I/ha]	Spray solution application rate [l/ha]	
	Arable crops						
*	Cereals	2	4-8 leaves	14-18	1	-	
	Cerears		tillering	25-29	1.5		
4	Dapasaad	2	4-8 leaves	14-18	1		
8	Rapeseed	2	beginning of stem elongation	30-31	1		
	Maize 1		4-6 leaves		2	200-300	
S	Potatoes	1	inter-row closure	31-39	2	-	
×	Sugar beets	1	4-б leaves	14-16	2		
•	Soybean	1	development of side shoots and the main shoot	21-49	1		
6	Legumes	1	stem elongation	30-39	1		
	Vegetable crops						
K	Bulb vegetables e.g. onion, leek	1-2	development of harvestable vegetative plant parts	41-45	1-1.5	_	
			development of harvestable vegetative plant parts	46-49	1.5		
Ď	Cucurbits e.g. pumpkin, zucchini, cucumber	1	leaf development	16-19	0.8		
Ģ	Brassica plants e.g. cabbage, cauliflower, broccoli	1-3	leaf development	14-16	0.5-1		
			rosette growth	31-39	1		
			development of harvestable vegetative plant parts	41-45	1		
	Root vegetables e.g. carrot, celery, beet	1	leaf development	14-16	1		
()?	Leaf vegetables e.g. lettuce, spinach	2	leaf development	14-19	0.5-1		
			development of harvestable vegetative plant parts	41-45	0.5-1		
2	Solanaceous e.g. tomato, pepper, early potato	1-4	leaf development and formation of side shoots	13-29	1		
			shoot elongation and tuber formation	31-49	1		
			inflorescence emergence, flowering, fruit development	51-79	1		
			ripening of fruit and seeds, withering	81-91	1		
) ?	Legumes	2-3	leaf development	13-15	0.5		
			leaf development	16-19	1		
	e.g. bean, pea		development of side shoots	21-29	1		
	Orchard crops						
7.	Stone-fruit trees e.g. sour cherry, sweet cherry	1-2	fruit development	72-79	0.3-0.6	- 500 900	
Ď	Pome trees e.g. apple, pear	1-2	fruit development	74-79	0.5-1	- 500-800	
0	Soft fruits e.g. strawberry, blueberry	1-2	beginning of vegetation, leaf development	10-19	0.2-0.4	300-500	



Przedsiębiorstwo Produkcyjno-Consultingowe ADOB Sp. z o.o. Sp. k.

ul. Kołodzieja 11 61-070 Poznań, PL email: office@adob.com.pl

